

Spiral Chains: The Proofs of Tait's and Tutte's Three-Edge-Coloring Conjectures

I. Cahit

Abstract

In this paper we have shown without assuming the four color theorem of planar graphs that every (bridgeless) cubic planar graph has a three-edge-coloring. This is an old-conjecture due to Tait in the sequel of efforts in settling the four-color conjecture at the end of the 19th century. We have also shown the applicability of our method to another well-known three edge-coloring conjecture on cubic graphs. Namely Tutte's conjecture that "every 2-connected cubic graph with no Petersen minor is 3-edge colorable". Hence the conclusion of this paper implies another non-computer proof of the four color theorem by using spiral-chains in different context.

To the memory of W.T. Tutte

1 Introduction

Assuming the four color theorem (4CT) it is easy to show that every cubic (bridgeless) planar graph has a three-edge-coloring since Tait found an equivalent formulation of the 4CT in terms of 3-edge-coloring.¹ This is an old-conjecture due to Tait in the sequel of efforts in settling the four-color conjecture at the end of the 19th century. In this paper we have given an independent proof of this famous conjecture. Again as we have recently introduced the new concept "spiral chains" in maximal planar graphs in the new non-computer proof of the four color theorem, we continue the use of spiral chains in the cubic plane graph for its three-edge-coloring [15]. Note that there are other equivalent formulation of 4CT in terms of three edge-coloring of the pairs of binary trees [4],[5]. We have first shown that any cubic planar graph can be decomposed into vertex disjoint spiral chains which in turn expressed as the union of *folded*²

¹Assume that the regions of a cubic planar graphs colored properly by red (R), blue (B), green (G) and yellow (Y) colors. Then if two neighbor regions are colored by RG or GY then color its boarder (edge in the cubic graph) by G, if two neighbor regions are colored by RG or BY then color it boarded by R and if two neighbor regions are colored by RY or BG then color its boarder by B. Then the theorem follows.

²Some of the edges are called "hairs" used twice, where *hair*-edge is the edge of the comb-caterpillar other than the backbone-edge (an edge having two ends degree three).

special spiral caterpillar (spiral-combs). Then based on the spiral-comb(s) decomposition we have provided a simple three-edge-coloring algorithm for any given cubic planar graph. In the second part of the paper we will show that the same method can also be applied to a similar but more general three-edge cubic graph (non-planar) coloring conjecture of Tutte which has been recently settled by a series lengthy papers using the classical four color theorem proof technique. That is

Conjecture 1 *Every 2-connected cubic graph with no Petersen minor is 3-edge colorable.*

This extends the four color theorem as Tait showed that the four color theorem is equivalent to the statement that every planar 2-connected cubic graph is 3-edge colorable. It also implies that certain non-planar graphs are 3-edge colorable. Let us say that G is *apex* if $G \setminus v$ is planar for some v and G is *doublecross* if it can be drawn in the plane with crossings, but with atmost two crossings and with all the crossings on the boundary of the finite region. Both apex and doublecross graphs have no Petersen minor so Tutte's conjecture implies:

Conjecture 2 (RST) (1) *Every 2-connected apex cubic graph is 3-edge colorable and (2) every 2-connected doublecross cubic graph is 3-edge colorable.*

The above conjecture is the outcome of joint efforts of Robertson, Seymour and Thomas (RST) [6,7,8] and it appears that excluding Roberstson, Sanders joins in the team for edge 3-coloring of cubic apex and doublecross graphs [9,10].

Tutte and RST equivalences has been shown in [6] that "a minimum counter-example" mean a 2-connected cubic graph G with no Petersen minor which is not 3-edge colorable, with $|V(G)|$ minimum. That is every minimum counter-example is either apex or doublecross. Eventually Tutte's conjecture has been settled by a series of papers [6]-[10] by using the classical proof method of the four color theorem [12],[13],[14]. The concept of spiral chain has been introduced by the author for an non-computer proof of the four-color theorem in 2004 [15]. We have applied spiral-chain coloring technique to the maximal planar graphs and we will use it the same for cubic planar and almost planar (apex and doublecross) cubic graphs. Therefore the main aim of this paper is to give an independent proof of the famous Tait's conjecture for bridgeless cubic planar graphs by using spiral-chain edge-coloring and then to show that it would be possible to extend it to the proof of conjecture RST. Hence to the proof of Conjecture 1.

2 Spiral-chains in cubic graphs

Throughout the paper the numbers 1, 2, 3 attached on the edges of a graph correspond, respectively to the colors (R)ed, (O)range and (G)reen³. That is $C = \{R, O, G\} \equiv \{1, 2, 3\}$. Let $G = G_0$ be a 2-connected (bridgeless) cubic planar graph drawn in the plane without crossings. Let B be denote its outer-boundary cycle consist of vertices $v_1, v_2, \dots, v_k, k \geq 3$. Without loss of generality select the edges starting from vertex v_1 ends at vertex v_k in the clockwise-direction (or anti-clockwise-direction) in the plane. That is we have selected the path from vertex v_1 to v_k on the B . We can do this always by selecting unused right-most edge in the plane, that is if the vertex v_i is in the spiral S then v_{i+1} will be in the S only if (v_i, v_{i+1}) is the right-most edge incident at v_i . Delete these selected vertices $v_1, v_2, \dots, v_k$ from the boundary cycle of G_0 and obtain the subgraph $G_1 = G_0 \setminus \{v_1, v_2, \dots, v_k\}$. Note that we have not selected the edge (v_k, v_1) since $(v_k, v_1) \cup P(v_1, v_k)$ is the outer-boundary cycle of G_0 . Then continue in this way for the other (inner) vertices on the outer-boundary vertices of the subgraph G_1 , that is select the edge (v_k, v_{k+1}) , where $v_k \in V(G_0)$ and $v_{k+1} \in V(G_1)$ and the outer-boundary vertices $v_{k+1}, v_{k+2}, \dots, v_{k+p}$ of G_1 in the clockwise direction and delete them from G_1 and the repeat the same process for $G_2 = G_1 \setminus \{v_{k+1}, v_{k+2}, \dots, v_{k+p}\}$. Let us assume that repetition of this spirally selection of the vertices results an empty graph at stage s . Now union of deleted vertices of $G_0, G_1, G_2, \dots, G_{s-1}$ form the spiral-chain S of G . We may describe spiral-chain S in terms of consecutive edges between the deleted edges as $\{e_1, e_2, e_3, \dots, e_{k-1}\} \cup (e_k) \cup \{e_{k+1}, e_{k+2}, \dots, e_{k+p-1}\} \cup \dots$, where $e_k = (v_{k-1}, v_{k+1})$ is the edge connecting two spiral segments of G_0 and G_1 . Not all 2-connected cubic planar graphs are hamiltonian, see for example famous Tutte graph in Fig.2. Otherwise the proof of Tait's conjecture would be immediately obtained by coloring hamiltonian cycle with colors 1 and 2 and coloring other edges with color 3. But deciding whether a cubic planar graph has an hamiltonian cycle or not is not an easy task. Therefore on Tait coloring of cubic planar graphs hamiltonicity is not our primary concern. On the other hand a 2-connected cubic planar graph may have a hamiltonian cycle but have several spiral chains (see Fig. 1 and 5). Edge coloring of these figures will be explained later.

Let T be tree with n vertices. Base tree T_b of T is obtained by removing end-vertices of T . A tree T is called *caterpillar* if its base tree is a path. We will use in our spiral-chain edge-coloring a special caterpillar called comb-tree:

Definition 3 *Comb tree T_c is a caterpillar in which all vertex degrees are 3 or 1.*

³These selection of colors may be further informational in the description of spiral-chain edge coloring in the cubic graph when they are viewed as the colors of the "traffic lights". That is Red for STOP, Orange for READY and Green for GO.

Figure 1: (a) Coloring by the use of an hamiltonian cycle, (b) spiral-chain edge-coloring

We will be investigating 3-edge coloring of 2-connected cubic planar graphs under two classes of cubic planar graphs since the "triangles" (cycle of length three) are the main issue in the uniqueness of the three colorings in planar graphs. For example T. Fowler and R. Thomas have settled a conjecture of Fiorini and Wilson, and Fisk about uniquely colorable planar graphs e.g.,

Theorem 4 *A 3-regular planar graph has a unique edge 3-coloring if and only if it can be obtained from the complete graph on four vertices by repeatedly replacing vertices by triangles.*

We will return to the uniquely edge 3-coloring planar graphs from the point-view of spiral-chain coloring.

3 Spiral-chain edge three-coloring

3.1 Triangle-free cubic bridgeless planar graphs

Let us start with a result on three edge-coloring of cubic planar graphs with triangles.

Theorem 5 *Three edge-coloring of a cubic planar graph G_Δ with triangles can be obtained from three edge-coloring of a triangle-free cubic bridgeless planar graph G .*

Proof. Let G_Δ be denote a cubic (bridgeless) planar graph. Then G_Δ can only contains (a) vertex disjoint triangles, (b) two triangles with only one edge common, call it *twin-triangle* and (c) if it contains three mutually adjacent triangles then $G_\Delta = K_4$. Assume that all triangle-free cubic planar graphs have been three edge-colored. Consider any cubic graph G_Δ which contains an triangle C_3 without a common edge with another triangle. That is let $V(C_3) = \{v_1, v_2, v_3\}$ and $E(C_3) = \{(v_1, v_2), (v_2, v_3), (v_3, v_1)\}$. G_Δ must have also vertices v_4, v_5, v_6 such that $(v_1, v_4), (v_2, v_5), (v_3, v_6) \in E(G_\Delta)$ where $v_4 \neq v_5 \neq v_6$. Let $C = \{i, j, k\}, i \neq j \neq k$ be the set of the three colors, i.e., $i, j, k \in \{1, 2, 3\}$. In any proper edge-coloring of G_Δ the subgraph formed by the triangle C_3 together with the edges $(v_1, v_4), (v_2, v_5), (v_3, v_6)$ must have an unique edge-coloring as shown in Fig. 2(a). Since the colors of the edges $(v_1, v_4), (v_2, v_5), (v_3, v_6)$ are i, j, k we can shrink the triangle C_3 into a vertex u_1 and obtain a new cubic graph (see also Fig. 2(a)). We repeat this process for every triangle till we arrive to a cubic-graph without distinct triangles. Order the new vertices so created as $u_1, u_2, \dots, u_k$. If G_Δ contains twin-triangle then it must have vertices v_1, v_2, v_3, v_4 and edges $(v_1, v_2), (v_2, v_3), (v_2, v_4), (v_1, v_3), (v_3, v_4)$ and two more edges (v_1, v_5) and (v_4, v_6) to make vertex degrees of the twin-triangle all three. Now all possible three edge-coloring of the subgraph has been shown in Fig. 2(b). Note that we have $v_5 \neq v_6$ otherwise it implies a bridge in G_Δ . Since the color of the edges (v_1, v_5) and (v_4, v_6) is $i \in \{1, 2, 3\}$ we delete the twin-triangle from G_Δ and join v_5 and v_6 with an new edge $e_1 = (v_5, v_6)$ to obtain another possibly triangle free cubic planar graph. If not we repeat the above processes for

Figure 2: Deletion of triangles from cubic planar graphs.

each twin-triangles in G_Δ till we get an triangle-free cubic planar graph G . Let $e_1, e_2, \dots, e_q$ be set of new edges inserted during the deletion of twin-triangles. Since we have assumed that G is a three edge-colorable we start from a coloring of G and re-insert all the deleted triangles for each $u_i \in V(G), i = 1, 2, \dots, k$ and twin-triangles for each $e_i \in E(G), i = 1, 2, \dots, q$ back into G but this time with suitable edge-colors based on the coloring of G and the cases in Fig. 2. Hence we have obtained three edge-coloring of G_Δ . Note that K_4 is an isolate case and is shown in Fig. 2(c). ■

3.2 Uniquely 3-edge colorable planar graphs

Characterization of uniquely 3-edge colorable graphs have been given by Towler and Thomas. Since these cubic planar graphs can only be obtained from the complete graph on four vertices by repeatedly replacing vertices by triangles, we note that they are hamiltonian and admit a very simple unique 3-edge coloring e.g., consider the coloring of Fig. 2(a). Let G_u be an uniquely 3-edge colorable graph. Let the vertices (v_i, v_j, v_k) induce an triangle in G_u and let $C = \{1, 2, 3\}$. We say the edge colors of an triangle is positive "+" if the colors of the edges $(v_i, v_j), (v_j, v_k), (v_k, v_i)$ respectively are 1, 2, 3 and negative "-" if the colors of its edges respectively are 2, 1, 3. That is in a positively signed triangle, numbers corresponding colors are rotated in clockwise direction while negatively signed triangle numbers are rotated in anti-clockwise direction (see Fig. 5). We also say that a vertex v in G_u is mature if it is on the corner of an triangle otherwise we call immature vertex. If all vertices of G_u are mature we say the graph is complete and denote it by K_u , i.e., complete with respect to the number of

triangles. For example the cubic graph shown in Fig. 5 is a complete graph on six triangles. Some properties of G_u are:

- Except K_4 (complete graph with four vertices), for any cubic graph G_u no two triangles have a common edge.
- Let S be a spiral-chain in G_u . For an 3-edge coloring of G_u the sign of all triangles incident to outer-boundary cycle have the same polarity, say all are $+$. Furthermore because of the character of the construction of uniquely 3-edge colorable graphs they constitute shell-structure (known also anti-matroids) of triangles in the graph and triangles in the consecutive shells must be labeled with opposite signs. For example in Fig. 5 all triangles labeled with positive signs are the triangles in the outer shell and neighbor shell triangles are labeled with negative signs. Since spiral-chains are shelling structure [15], we can use this property when we color the edges of G_u . That is we can extract the sign of the triangle from the knowledge of the location of the edge in the spiral chain and from the sign of the previous triangle when we meet one along the edges of the spiral chain. We omit details of the spiral-chain coloring algorithm for 3-edge uniquely colorable graphs since the algorithm given in the next section may as well be applied for this class of cubic planar graphs.

3.3 Spiral-chain edge-coloring

In this section we will give an three edge-coloring algorithm which uses spiral chains of the given cubic bridgeless planar graph. Let $C = \{R, O, G\}$. In the figures we use numbers 1, 2, 3 on the edges instead of letters to avoid confusion. Let S be denote a spiral-chain of G . As we go through the vertices of S we construct a path P_s with vertices $v_1, v_2, \dots, v_k$ and assume that $|V(P)| = |V(G)|$, i.e., $k = n$. We will investigate cubic graphs with several spiral-chains later. Edges incident to P_s together with the edges $E(P_s)$ of the path constitute a comb-caterpillar tree $C_s(V_s, E_s)$, where $V_s = \{v_1, v_2, \dots, v_k\}$ and $E_s = E(P_s) \cup \{\overline{E(P_s)} : \{v_i, v_j\} \mid v_i, v_j \in V(P_s) \text{ and } d(v_i, v_j) \geq 2\}$. An edge b of the C_s is called the *backbone*-edge if $b \in E(P_s)$ and is called the *hair*-edge h if $h \in \overline{E(P_s)}$. It is clear that $C_s = G$ but some of the edges of $\overline{E(P_s)}$ are counted twice since any hair-edge of C_s has two ends in the spiral-chain S . Again as we go through the edges of S we call that hair edge $h \in \overline{E(P_s)}$ is *real* if it is encountered for the first-time and *imaginary* if it is encountered for the second-time. In the figures all the real-hair edges have been shown in straight-lines while the imaginary-hair edges are shown with dashed-lines. The importance of the real and imaginary hairs will be clearer when deciding the color of the current edge in the spiral chain. We have classify the colors into two classes. The color green G is called as the "primary" color while the colors orange O and red R are called the "secondary" colors. In the coloring algorithm given below primary color G is used mostly for the spiral-chain backbone edges and the secondary colors O and R are used mostly for the hair-edges.

Algorithm A. *Spiral-chain edge-coloring (for single spiral-chain)*

Let the cubic planar graph G be drawn in plane without any crossings. As the first step construct in clockwise direction a spiral-chain S of G starting from a vertex v_1 of its outer-boundary cycle. Let C_s be the corresponding comb of S . Start coloring the edges of the backbone edges of C_s alternately by green and orange colors i.e., (G, O) -chain and the hair edges of C_s by the red R color. If all vertices of S is covered and edges incident to vertices are properly colored by G, O , and R then the coloring is the desired proper three edge-coloring.

a) If for some backbone edge $e_i = (v_i, v_{i-1})$ is forced to be colored by R because of the color (it must be orange O since G is primary color) of the hair h_{i-1} at v_{i-1} . Furthermore if the hair h_i had been colored by R before then both e_i and h_i would be R . In this case there must be a vertex v_j , with $j > i$ and a spiral sub-path $P(v_r, v_s)$, with $s > r < j > i$. But the path $P(v_i, v_j) = (v_i, v_r) \cup P(v_r, v_s) \cup (v_s, v_j)$ is a (R, O) -Kempe chain, where (v_i, v_r) and (v_s, v_j) are the hair-edges. Therefore we perform Kempe-chain color switching for the edges of $P(v_i, v_j)$ and resolves the problem at vertex v_i .

a) If for some backbone edge $e_i = (v_i, v_{i-1})$ is forced to be colored by O because of the color of the hair h_{i-1} at v_{i-1} . Furthermore if the hair h_i had been colored by O before then both e_i and h_i would be O . In this case there must be a vertex v_j , with $j > i$ and a spiral sub-path $P(v_r, v_s)$, with $s > r < j > i$. But the path $P(v_i, v_j) = (v_i, v_r) \cup P(v_r, v_s) \cup (v_s, v_j)$ is a (O, G) -Kempe chain, where (v_i, v_r) and (v_s, v_j) are the hair-edges. Therefore we perform Kempe-chain color switching for the edges of $P(v_i, v_j)$ and resolves the problem at vertex v_i .

That is in critical cases we have always suitable Kempe-chain with both ends are hairs of the spiral comb-tree. Hence we can perform Kempe-chain color switching within the spiral-chain of G when we needed to resolve color duplication at vertex v_i .

Fig. 1(b) and Fig. 3(b) illustrate the spiral-chain edge-coloring algorithm.

Algorithm A can be easily modified to cubic planar graphs having more than one spiral chains. In this case if the current vertex, say v_a of spiral-chain S_i adjacent to two vertices v_b, v_c that have already been contained in S_i before or in some other spiral-chain $S_j, j < i$ then we cannot go further. Hence we select closest new vertex incident $S_k, k = 1, 2, \dots, i$ and start new spiral-chain S_{i+1} from that vertex. When all vertices of G covered we obtain vertex disjoint set of spiral-chains $S_1, S_2, \dots, S_m$. Note that some of spiral-chains degenerate to a single vertex. One can easily notice that spiral-chains $S_1, S_2, \dots, S_m$ are topologically nested in the plane. See for example spiral-chains in Fig. 5 and 6. Now we ready to give spiral-chain edge-coloring algorithm for this case:

Algorithm B. Spiral-chain edge-coloring (for multiple spiral-chains).

Let $S_1, S_2, \dots, S_m$ be the set of spiral-chains of the cubic planar graph. Apply Algorithm A to each spiral-chain $S_i, i = 1, 2, \dots, m$. Note that for each application of Algorithm A we select always the color "green" G as the primary color. This selection would minimize execution of number of the steps (a) and (b) in Algorithm A. That is it will color in green, maximum number of the backbone edges in the spiral-chains.

Figure 3: (a) Spiral-chain and spiral-cut, (b) and its spiral three edge-coloring.

Figure 4: Spiral comb tree in the coloring algorithm.

4 Tutte's three edge-coloring conjecture

The main idea of this section is to show that possible cubic graph minimal counter-examples in Conjecture 2 (RST) can be three edge-colorable by using spiral-chain edge coloring algorithm that has been given in the previous section. In other words the contribution of this section is a finishing touch to the deep result of Robertson, Seymour and Thomas [6] that the only possible minimal counter-examples to Tutte's conjecture (Conjecture 1) would be apex and doublecross cubic non-planar graphs. We only need to give a suitable modification in the construction of spiral chains since the possible counter-examples are specific non-planar cubic graphs i.e., apex and doublecross. Let G_{ad} denote apex or doublecross cubic graph.

Let us note the following observation:

In the previous section we have agreed that spiral-chains in the cubic planar graphs are rotated in the clockwise or anti-clockwise. One of the reason of this is to *not* loss our way in the edge-coloring algorithm in the cubic planar graph and ordering the spiral-chains in an nested fashion. In any cubic non-planar graph G_{ad} planarity are destroyed either by a special vertex or two crossings of a pair of edges within the same face. Suppose we start construction of the spiral-chain in clockwise direction in G_{ad} we see that for each edge-crossing the direction of spiral-chain is switched from clockwise to anti-clockwise and *vice versa*. Therefore spiral-chain edge-coloring algorithm described in the previous section can be applied without any difficulty to the graph G_{ad} .

For example illustration of apex and doublecross graphs are taken from [11] and have been shown with the spiral-chain edge colorings.

5 Conclusion

In this paper we have shown the effectiveness of spiral-chain coloring in the proofs of the well-known Tait's and Tutte's three edge-coloring of cubic graphs. With this result we have also given in a way the proof of the four color theorem in two-ways either by using spiral-chain vertex-coloring in the maximal planar or by using spiral-chain edge-coloring for bridgless cubic planar graphs. We hope to announce similar results in an near future for the Steinberg's vertex three-coloring conjecture of planar graphs and for the Hadwiger's conjecture which is a generalization of the four color problem.

Figure 5: Spiral-chain coloring of an uniquely edge 3-colorable cubic graph.

Figure 6: Spiral-chain coloring of the Tutte graph.

Figure 7: An apex and its spiral-chain coloring.


Figure 8: A doublecross and its spiral-chain coloring.

References


- [1] P.G. Tait, "On the colouring of maps", *Proceedings of the Royal Society of Edinburgh Section A*, 10: 501-503, 1878-1880.
- [2] P.G. Tait, "Remarks on the previous communication", *Proceedings of the Royal Society of Edinburgh Section A*, 10:729, 1878-1880.
- [3] P.G. Tait, "Note on a theorem in geometry of position", *Transactions of the Royal Society of Edinburgh*, 29:657-660, 1880.
- [4] A. Czumaj and A. Gibbons, "Guthrie's problem: new equivalences and rapid reductions", *Theoretical Computer Science*, 154:3-22, 1996.
- [5] P. Sant, "Algorithmics of edge-colouring pairs of 3-regular trees", Ph.D. thesis, University of London, December 2003.
- [6] N. Robertson, P.D. Seymour and R. Thomas, "Tutte's edge-colouring conjecture", *J. Combin. Theory Ser. B* 70 (1997), 166-183.
- [7] N. Robertson, P.D. Seymour and R. Thomas, "Cyclically 5-connected cubic graphs", in preparation.
- [8] N. Robertson, P.D. Seymour and R. Thomas, "Excluded minors in cubic graphs", in preparation.
- [9] D. P. Sanders and R. Thomas, "Edge 3-coloring cubic apex graphs", in preparation.
- [10] D. P. Sanders, P.D. Seymour and R. Thomas, "Edge 3-coloring cubic doublecross graphs", in preparation.
- [11] W.T. Tutte, "On algebraic theory of graph coloring", *J. Combin. Theory*, 1 (1966), 15-50.
- [12] R. Thomas, "An update on the four-color theorem", *Notices Amer. Math. Soc.*, 45 (1998), 7, 848-859.
- [13] K. Appel and W. Haken, "Every planar map is four colorable", *Contemporary Math.*, 98 (1989).
- [14] N. Robertson, D. P. Sanders, P. D. Seymour and R. Thomas, "The four colour theorem", *J. Combin. Theory Ser. B* 70 (1997), 2-44.
- [15] I. Cahit, "Spiral chains: A new proof of the four color theorem", arXiv preprint, math.CO/0408247, August 18, 2004.

I. Cahit
 Near East University, North Cyprus
 E-mail: icahit@ebim.com.tr or icahit@gmail.com

July 6, 2005




(b)




(Orange-Green)-Kempe chain

(Red-Orange)-Kempe chain




(a)




STEP 1: (Spira-chain construction)




STEP 2: (Spira-chain coloring)




(a)





(b)




(c)









○ — ○ : Edge encountered first-time { — } : Green is the *primary* color
 ○ - - - ○ : Edge encountered second-time { — , — } : Red and orange are the *secondary* colors

